



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>A61K 38/29, C07K 1/06, 14/635</b>	<b>A1</b>	(11) International Publication Number: <b>WO 97/02834</b> (43) International Publication Date: 30 January 1997 (30.01.97)								
<p>(21) International Application Number: <b>PCT/US96/11292</b></p> <p>(22) International Filing Date: <b>3 July 1996 (03.07.96)</b></p> <p>(30) Priority Data:</p> <table border="0"> <tr> <td>60/001,105</td> <td>13 July 1995 (13.07.95)</td> <td>US</td> </tr> <tr> <td>60/003,305</td> <td>6 September 1995 (06.09.95)</td> <td>US</td> </tr> <tr> <td>08/626,186</td> <td>29 March 1996 (29.03.96)</td> <td>US</td> </tr> </table> <p>(60) Parent Application or Grant (63) Related by Continuation US Filed on <b>08/626,186 (CIP)</b> <b>29 March 1996 (29.03.96)</b></p> <p>(71) Applicant (for all designated States except US): <b>BIOMEASURE INCORPORATED [US/US]; 27 Maple Street, Milford, MA 01757 (US).</b></p> <p>(72) Inventor; and (75) Inventor/Applicant (for US only): <b>DONG, Zheng, Xin [-/US]; 40 Angelica Drive, Framingham, MA 01701 (US).</b></p> <p>(74) Agent: <b>TSAO, Y., Rocky; Fish &amp; Richardson P.C., 225 Franklin Street, Boston, MA 02110 (US).</b></p>	60/001,105	13 July 1995 (13.07.95)	US	60/003,305	6 September 1995 (06.09.95)	US	08/626,186	29 March 1996 (29.03.96)	US	<p>(81) Designated States: <b>AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</b></p> <p><b>Published</b> <i>With international search report.</i></p>
60/001,105	13 July 1995 (13.07.95)	US								
60/003,305	6 September 1995 (06.09.95)	US								
08/626,186	29 March 1996 (29.03.96)	US								
<p>(54) Title: <b>ANALOGS OF PARATHYROID HORMONE</b></p> <p>(57) Abstract</p> <p>Peptide variants of fragment (1-34) of parathyroid hormone, in which at least one of the amino acid residues at positions 7, 11, 23, 24, 27, 28 and 31 is cyclohexylalanine, or at least one of the amino acid residues at positions 3, 16, 17, 18, 19 and 34 is <math>\alpha</math>-aminoisobutyric acid; or, alternatively, at least the amino acid residue at position 1 is <math>\alpha,\beta</math>-diaminopropionic acid, the amino acid residue at position 27 is homoarginine, or the amino acid residue at position 31 is norleucine.</p>										

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

- 1 -

## ANALOGS OF PARATHYROID HORMONE

### Background of the Invention


Parathyroid hormone ("PTH") is a polypeptide  
5 produced by the parathyroid glands. The mature  
circulating form of the hormone is comprised of 84 amino  
acid residues. The biological action of PTH can be  
reproduced by a peptide fragment of its N-terminus (e.g.  
amino acid residues 1 through 34). Parathyroid hormone-  
10 related protein ("PTHrP") is a 139 to 173 amino  
acid-protein with N-terminal homology to PTH. PTHrP  
shares many of the biological effects of PTH including  
binding to a common PTH/PTHrP receptor. Tregear, et al.,  
Endocrinol., 93:1349 (1983). PTH peptides from many  
15 different sources, e.g., human, bovine, rat, chicken,  
have been characterized. Nissenson, et al., Receptor,  
3:193 (1993).

PTH has been shown to both improve bone mass and  
quality. Dempster, et al., Endocrine Rev., 14:690  
20 (1993); and Riggs, Amer. J. Med., 91 (Suppl. 5B):37S  
(1991). The anabolic effect of intermittently  
administered PTH has been observed in osteoporotic men  
and women either with or without concurrent  
antiresorptive therapy. Slovik, et al., J. Bone Miner.  
25 Res., 1:377 (1986); Reeve, et al., Br. Med. J., 301:314  
(1990); and Hesch, R-D., et al., Calcif. Tissue Int'l,  
44:176 (1989).

### Summary of the Invention

In one aspect, the invention relates to peptide  
30 variants of PTH(1-34) of the following generic formula:

- 2 -



wherein

- 10            $A_1$  is Ser, Ala, or Dap;  
               $A_3$  is Ser, Thr, or Aib;  
               $A_5$  is Leu, Nle, Ile, Cha,  $\beta$ -Nal, Trp, Pal, Phe or  
              p-X-Phe, in which X is OH, a halogen, or  $\text{CH}_3$ ;  
               $A_7$  is Leu, Nle, Ile, Cha,  $\beta$ -Nal, Trp, Pal, Phe, or  
              p-X-Phe in which X is OH, a halogen, or  $\text{CH}_3$ ;  
 15            $A_8$  is Met, Nva, Leu, Val, Ile, Cha, or Nle;  
               $A_{11}$  is Leu, Nle, Ile, Cha,  $\beta$ -Nal, Trp, Pal, Phe or  
              p-X-Phe in which X is OH, a halogen, or  $\text{CH}_3$ ;  
               $A_{12}$  is Gly or Aib;  
               $A_{15}$  is Leu, Nle, Ile, Cha,  $\beta$ -Nal, Trp, Pal, Phe,  
 20 or p-X-Phe in which X is OH, a halogen, or  $\text{CH}_3$ ;  
               $A_{16}$  is Ser, Asn, Ala, or Aib;  
               $A_{17}$  is Ser, Thr, or Aib;  
               $A_{18}$  is Met, Nva, Leu, Val, Ile, Nle, Cha, or Aib;  
               $A_{19}$  is Glu or Aib;  
 25            $A_{21}$  is Val, Cha, or Met;  
               $A_{23}$  is Trp or Cha;  
               $A_{24}$  is Leu or Cha;  
               $A_{27}$  is Lys, Aib, Leu, hArg, Gln, or Cha;  
               $A_{28}$  is Leu or Cha;  
 30            $A_{30}$  is Asp or Lys;  
               $A_{31}$  is Val, Nle, Cha, or deleted;  
               $A_{32}$  is His or deleted;  
               $A_{33}$  is Asn or deleted;  
               $A_{34}$  is Phe, Tyr, Amp, Aib, or deleted;

- 3 -

each of  $R_1$  and  $R_2$  is, independently, H,  $C_{1-12}$  alkyl,  $C_{2-12}$  alkenyl,  $C_{7-20}$  phenylalkyl,  $C_{11-20}$  naphthylalkyl,  $C_{1-12}$  hydroxyalkyl,  $C_{2-12}$  hydroxyalkenyl,  $C_{7-20}$  hydroxyphenylalkyl, or  $C_{11-20}$  hydroxynaphthylalkyl; or  
5 one and only one of  $R_1$  and  $R_2$  is  $COE_1$  in which  $E_1$  is  $C_{1-12}$  alkyl,  $C_{2-12}$  alkenyl,  $C_{7-20}$  phenylalkyl,  $C_{11-20}$  naphthylalkyl,  $C_{1-12}$  hydroxyalkyl,  $C_{2-12}$  hydroxyalkenyl,  $C_{7-20}$  hydroxy-phenylalkyl, or  $C_{11-20}$  hydroxynaphthylalkyl; and  
 $R_3$  is OH,  $NH_2$ ,  $C_{1-12}$  alkoxy, or  $NH-Y-CH_2-Z$  in which  
10 Y is a  $C_{1-12}$  hydrocarbon moiety and Z is H, OH,  $CO_2H$ , or  $CONH_2$ ;

provided that (i) at least one of  $A_5$ ,  $A_7$ ,  $A_8$ ,  $A_{11}$ ,  $A_{15}$ ,  $A_{18}$ ,  $A_{21}$ ,  $A_{23}$ ,  $A_{24}$ ,  $A_{27}$ ,  $A_{28}$ , and  $A_{31}$  is Cha, or at least one of  $A_3$ ,  $A_{12}$ ,  $A_{16}$ ,  $A_{17}$ ,  $A_{18}$ ,  $A_{19}$ , and  $A_{34}$  is Aib;  
15 or that (ii) at least  $A_1$  is Dap,  $A_7$  is  $\beta$ -Nal, Trp, Pal, Phe, or p-X-Phe,  $A_{15}$  is  $\beta$ -Nal, Trp, Pal, Phe, or p-X-Phe,  $A_{27}$  is hArg, or  $A_{31}$  is Nle; or a pharmaceutically acceptable salt thereof.

A subset of the compounds covered by the above  
20 formula are those in which at least one of  $A_5$ ,  $A_7$ ,  $A_{11}$ ,  $A_{15}$ ,  $A_{18}$ ,  $A_{21}$ ,  $A_{23}$ ,  $A_{24}$ ,  $A_{27}$ ,  $A_{28}$ , and  $A_{31}$  is Cha. For example,  $A_3$  is Ser;  $A_5$  is Ile;  $A_7$  is Leu or Cha;  $A_8$  is Met, Nva, Leu, Val, Ile, or Nle;  $A_{11}$  is Leu or Cha;  $A_{12}$  is Gly;  $A_{15}$  is Leu or Cha;  $A_{16}$  is Asn or Aib;  $A_{17}$  is Ser;  $A_{18}$   
25 is Met or Nle;  $A_{21}$  is Val;  $A_{27}$  is Lys, hArg, or Cha;  $A_{32}$  is His;  $A_{31}$  is Val, Nle, or Cha;  $A_{33}$  is Asn;  $A_{34}$  is Phe, Tyr, Amp, or Aib;  $R_1$  is H;  $R_2$  is H; and  $R_3$  is  $NH_2$ ;  
provided that at least one of  $A_5$ ,  $A_7$ ,  $A_8$ ,  $A_{11}$ ,  $A_{15}$ ,  $A_{18}$ ,  $A_{21}$ ,  $A_{23}$ ,  $A_{24}$ ,  $A_{27}$ ,  $A_{28}$ , and  $A_{31}$  is Cha, or at least one of  
30  $A_3$ ,  $A_{12}$ ,  $A_{16}$ ,  $A_{17}$ ,  $A_{18}$ ,  $A_{19}$ , and  $A_{34}$  is Aib. If desired, at least one of  $A_7$  and  $A_{11}$  can be Cha; or at least one of  $A_{15}$ ,  $A_{23}$ ,  $A_{24}$ ,  $A_{27}$ ,  $A_{28}$ , and  $A_{31}$  is Cha.

In another subset, at least one of  $A_3$ ,  $A_{12}$ ,  $A_{16}$ ,  $A_{17}$ ,  $A_{18}$ ,  $A_{19}$ , and  $A_{34}$  is Aib. For example,  $A_3$  is Ser or  
35 Aib;  $A_5$  is Ile;  $A_7$  is Leu or Cha;  $A_8$  is Met, Nva, Leu,

- 4 -

Val, Ile, or Nle; A<sub>11</sub> is Leu or Cha; A<sub>15</sub> is Leu or Cha, A<sub>16</sub> is Asn or Aib; A<sub>18</sub> is Met, Aib, or Nle; A<sub>21</sub> is Val; A<sub>27</sub> is Lys, Aib, Leu, hArg, or Cha; A<sub>31</sub> is Val, Nle, or Cha; A<sub>32</sub> is His; A<sub>33</sub> is Asn; A<sub>34</sub> is Phe, Tyr, Amp, or Aib; R<sub>1</sub> is H; R<sub>2</sub> is H; and R<sub>3</sub> is NH<sub>2</sub>; provided that at least one of A<sub>5</sub>, A<sub>7</sub>, A<sub>8</sub>, A<sub>11</sub>, A<sub>15</sub>, A<sub>18</sub>, A<sub>21</sub>, A<sub>23</sub>, A<sub>24</sub>, A<sub>27</sub>, A<sub>28</sub>, and A<sub>31</sub> is Cha, or at least one of A<sub>3</sub>, A<sub>12</sub>, A<sub>16</sub>, A<sub>17</sub>, A<sub>18</sub>, A<sub>19</sub>, and A<sub>34</sub> is Aib. If desired, at least one of A<sub>7</sub> and A<sub>11</sub> can be Cha; or at least one of A<sub>15</sub>, A<sub>23</sub>, A<sub>24</sub>, A<sub>27</sub>, A<sub>28</sub>, and A<sub>31</sub> is Cha.

In a still further subset, at least one of A<sub>5</sub>, A<sub>7</sub>, A<sub>8</sub>, A<sub>11</sub>, A<sub>15</sub>, A<sub>18</sub>, A<sub>21</sub>, A<sub>23</sub>, A<sub>24</sub>, A<sub>27</sub>, A<sub>28</sub>, and A<sub>31</sub> is Cha, or at least one of A<sub>3</sub>, A<sub>12</sub>, A<sub>16</sub>, A<sub>17</sub>, A<sub>18</sub>, A<sub>19</sub>, and A<sub>34</sub> is Aib. For example, A<sub>3</sub> is Ser or Aib; A<sub>5</sub> is Ile; A<sub>7</sub> is Leu or Cha; A<sub>8</sub> is Met, Nva, Leu, Val, Ile, or Nle; A<sub>11</sub> is Leu or Cha; A<sub>15</sub> is Leu or Cha; A<sub>16</sub> is Asn or Aib; A<sub>18</sub> is Met, Aib, or Nle; A<sub>21</sub> is Val; A<sub>27</sub> is Lys, Aib, Leu, hArg, or Cha; A<sub>31</sub> is Val, Nle, or Cha; A<sub>32</sub> is His; A<sub>33</sub> is Asn; A<sub>34</sub> is Phe, Tye, Amp, or Aib; R<sub>1</sub> is H; R<sub>2</sub> is H; and R<sub>3</sub> is NH<sub>2</sub>. If desired, at least one of A<sub>7</sub> and A<sub>11</sub> is Cha and at least one of A<sub>16</sub>, A<sub>19</sub>, and A<sub>34</sub> is Aib; or at least one of A<sub>24</sub>, A<sub>28</sub>, and A<sub>31</sub> is Cha and at least one of A<sub>16</sub> and A<sub>17</sub> is Aib.

In yet another subset, at least one of A<sub>1</sub> is Dap, A<sub>7</sub> is  $\beta$ -Nal, Trp, Pal, Phe or p-X-Phe, A<sub>13</sub> is  $\beta$ -Nal, Trp, Pal, Phe, or p-X-Phe. For example, A<sub>1</sub> is Ser, Gly, or Dap; A<sub>3</sub> is Ser or Aib; A<sub>8</sub> is Met, Nva, Leu, Val, Ile, or Nle; A<sub>16</sub> is Asn or Aib; A<sub>18</sub> is Met, Aib, or Nle; A<sub>21</sub> is Val; A<sub>27</sub> is Lys, Aib, Leu, hArg, or Cha; A<sub>31</sub> is Val, Nle, or Cha; A<sub>32</sub> is His; A<sub>33</sub> is Asn; A<sub>34</sub> is Phe, Tyr, Amp, or Aib; R<sub>1</sub> is H; R<sub>2</sub> is H; and R<sub>3</sub> is NH<sub>2</sub>.


The following are examples of the peptide of this invention as covered by the above formula: [Cha<sup>7</sup>]hPTH(1-34)NH<sub>2</sub>; [Cha<sup>11</sup>]hPTH(1-34)NH<sub>2</sub>; [Cha<sup>15</sup>]hPTH(1-34)NH<sub>2</sub>; [Cha<sup>7, 11</sup>]hPTH(1-34)NH<sub>2</sub>; [Cha<sup>7, 11</sup>, Nle<sup>8, 18</sup>, Tyr<sup>34</sup>]hPTH(1-34)NH<sub>2</sub>; [Cha<sup>23</sup>]hPTH(1-34)NH<sub>2</sub>; [Cha<sup>24</sup>]hPTH(1-34)NH<sub>2</sub>; [Nle<sup>8, 18</sup>,

- 5 -

Cha<sup>27</sup>]hPTH(1-34)NH<sub>2</sub>; [Cha<sup>28</sup>]hPTH(1-34)NH<sub>2</sub>; [Cha<sup>31</sup>]hPTH(1-  
 34)NH<sub>2</sub>; [Cha<sup>27</sup>]hPTH(1-34)NH<sub>2</sub>; [Cha<sup>27, 29</sup>]hPTH(1-34)NH<sub>2</sub>;  
 [Cha<sup>28</sup>]bPTH(1-34)NH<sub>2</sub>; [Cha<sup>28</sup>]rPTH(1-34)NH<sub>2</sub>; [Cha<sup>24, 28,</sup>  
 31]hPTH(1-34)NH<sub>2</sub>; [Aib<sup>16</sup>]hPTH(1-34)NH<sub>2</sub>; [Aib<sup>19</sup>]hPTH(1-  
 5 34)NH<sub>2</sub>; [Aib<sup>34</sup>]hPTH(1-34)NH<sub>2</sub>; [Aib<sup>16, 19</sup>]hPTH(1-34)NH<sub>2</sub>;  
 [Aib<sup>16, 19, 34</sup>]bPTH(1-34)NH<sub>2</sub>; [Aib<sup>16, 34</sup>]hPTH(1-34)NH<sub>2</sub>;  
 [Aib<sup>19, 34</sup>]hPTH(1-34)NH<sub>2</sub>; [Cha<sup>7, 11, Nle<sup>8, 18, Aib<sup>16, 19,</sup></sup>  
 Tyr<sup>34</sup>]hPTH(1-34)NH<sub>2</sub>; [Cha<sup>7, 11, Nle<sup>8, 18, 31, Aib<sup>16, 19,</sup></sup>  
 Tyr<sup>34</sup>]hPTH(1-34)NH<sub>2</sub>; [Cha<sup>7, Aib<sup>16</sup></sup>]hPTH(1-34)NH<sub>2</sub>; [Cha<sup>11,</sup>  
 10 Aib<sup>16</sup>]hPTH(1-34)<sub>2</sub>; [Cha<sup>7, Aib<sup>34</sup></sup>]hPTH(1-34)NH<sub>2</sub>; [Cha<sup>11,</sup>  
 Aib<sup>34</sup>]hPTH(1-34)NH<sub>2</sub>; [Cha<sup>27, Aib<sup>16</sup></sup>]hPTH(1-34)NH<sub>2</sub>; [Cha<sup>27,</sup>  
 Aib<sup>34</sup>]hPTH(1-34)NH<sub>2</sub>; [Cha<sup>28, Aib<sup>16</sup></sup>]hPTH(1-34)NH<sub>2</sub>; [Cha<sup>28,</sup>  
 Aib<sup>34</sup>]hPTH(1-34)NH<sub>2</sub>; [Nle<sup>31</sup>]hPTH(1-34)NH<sub>2</sub>; [hArg<sup>27</sup>]hPTH(1-  
 34)NH<sub>2</sub>; [Dap<sup>1, Nle<sup>8, 18, Tyr<sup>34</sup></sup>]hPTH(1-34)NH<sub>2</sub>;  
 15 [Nle<sup>31</sup>]bPTH(1-34)NH<sub>2</sub>; [Nle<sup>31</sup>]rPTH(1-34)NH<sub>2</sub>; [hArg<sup>27</sup>]bPTH(1-  
 34)NH<sub>2</sub>; [hArg<sup>27</sup>]rPTH(1-34)NH<sub>2</sub>; [Cha<sup>7, 11, Aib<sup>19,</sup></sup>  
 Lys<sup>30</sup>]hPTH(1-34)NH<sub>2</sub>; [Aib<sup>12</sup>]hPTH(1-34)NH<sub>2</sub>; [Cha<sup>24, 28, 31,</sup>  
 Lys<sup>30</sup>]hPTH(1-34)NH<sub>2</sub>; [Cha<sup>28, 31</sup>]hPTH(1-34)NH<sub>2</sub>; [Cha<sup>7, 11,</sup>  
 Nle<sup>8, 18, Aib<sup>34</sup></sup>]hPTH(1-34)NH<sub>2</sub>; [Aib<sup>3</sup>]hPTH(1-34)NH<sub>2</sub>;  
 20 [Cha<sup>8</sup>]hPTH(1-34)NH<sub>2</sub>; [Cha<sup>15</sup>]hPTH(1-34)NH<sub>2</sub>; [Cha<sup>7, 11,</sup>  
 Aib<sup>19</sup>]hPTH(1-34)NH<sub>2</sub>; [Cha<sup>7, 11, Aib<sup>16</sup></sup>]hPTH(1-34)NH<sub>2</sub>;  
 [Aib<sup>17</sup>]hPTH(1-34)NH<sub>2</sub>; [Cha<sup>5</sup>]hPTH(1-34)NH<sub>2</sub>; [Cha<sup>7, 11,</sup>  
 15]hPTH(1-34)NH<sub>2</sub>; [Cha<sup>7, 11, Nle<sup>8, 18, Aib<sup>19,</sup></sup>  
 Tyr<sup>34</sup>]hPTH(1-34)NH<sub>2</sub>; [Cha<sup>7, 11, Nle<sup>8, 18, Aib<sup>19, Lys<sup>30,</sup></sup></sup>  
 25 Tyr<sup>34</sup>]hPTH(1-34)NH<sub>2</sub>; [Cha<sup>7, 11, 15</sup>]hPTH(1-34)NH<sub>2</sub>;  
 [Aib<sup>17</sup>]hPTH(1-34)NH<sub>2</sub>; [Cha<sup>7, 11, Leu<sup>27</sup></sup>]hPTH(1-34)NH<sub>2</sub>;  
 [Cha<sup>7, 11, 15, Leu<sup>27</sup></sup>]hPTH(1-34)NH<sub>2</sub>; [Cha<sup>7, 11, 27</sup>]hPTH(1-  
 34)NH<sub>2</sub>; [Cha<sup>7, 11, 15, 27</sup>]hPTH(1-34)NH<sub>2</sub>; [Trp<sup>15</sup>]hPTH(1-  
 34)NH<sub>2</sub>; [Nal<sup>15</sup>]hPTH(1-34)NH<sub>2</sub>; [Trp<sup>15, Cha<sup>23</sup></sup>]hPTH(1-34)NH<sub>2</sub>;  
 30 [Cha<sup>15, 23</sup>]hPTH(1-34)NH<sub>2</sub>; [Phe<sup>7, 11</sup>]hPTH(1-34)NH<sub>2</sub>; [Nal<sup>7,</sup>  
 11]hPTH(1-34)NH<sub>2</sub>; [Trp<sup>7, 11</sup>]hPTH(1-34)NH<sub>2</sub>; [Phe<sup>7, 11,</sup>  
 15]hPTH(1-34)NH<sub>2</sub>; [Nal<sup>7, 11, 15</sup>]hPTH(1-34)NH<sub>2</sub>; [Trp<sup>7, 11,</sup>  
 15]hPTH(1-34)NH<sub>2</sub>; and [Tyr<sup>7, 11, 15</sup>]hPTH(1-34)NH<sub>2</sub>..</sup></sup></sup></sup></sup>

In another aspect, this invention relates to  
 35 peptides covered by the following formula:

- 6 -



wherein

- $$\begin{array}{l}
 A_1 \text{ is Ala, Ser, or Dap;} \\
 A_3 \text{ is Ser or Aib;} \\
 A_5 \text{ is His, Ile, or Cha;} \\
 A_7 \text{ is Leu, Cha, Nle, } \beta\text{-Nal, Trp, Pal,} \\
 \text{Phe, or p-X-Phe in which X is OH, a halogen, or CH}_3\text{;} \\
 A_8 \text{ is Leu, Met, or Cha;} \\
 A_{10} \text{ is Asp or Asn;} \\
 A_{11} \text{ is Lys, Leu, Cha, Phe, or } \beta\text{-Nal;} \\
 A_{12} \text{ is Gly or Aib;} \\
 A_{14} \text{ is Ser or His;} \\
 A_{15} \text{ is Ile, or Cha;} \\
 A_{16} \text{ is Gln or Aib;} \\
 A_{17} \text{ is Asp or Aib;} \\
 A_{18} \text{ is Leu, Aib, or Cha;} \\
 A_{19} \text{ is Arg or Aib;} \\
 A_{22} \text{ is Phe, Glu, Aib, or Cha;} \\
 A_{23} \text{ is Phe, Leu, Lys, or Cha;} \\
 A_{24} \text{ is Leu, Lys, or Cha;} \\
 A_{25} \text{ is His, Aib, or Glu;} \\
 A_{26} \text{ is His, Aib, or Lys;} \\
 A_{27} \text{ is Leu, Lys, or Cha;} \\
 A_{28} \text{ is Ile, Leu, Lys, or Cha;} \\
 A_{29} \text{ is Ala, Glu, or Aib;} \\
 A_{30} \text{ is Glu, Cha, Aib, or Lys;} \\
 A_{31} \text{ is Ile, Leu, Cha, Lys, or deleted;} \\
 A_{32} \text{ is His or deleted;} \\
 A_{33} \text{ is Thr or deleted;}
 \end{array}$$



- 7 -

A<sub>34</sub> is Ala or deleted;

each of R<sub>1</sub> and R<sub>2</sub> is, independently, H, C<sub>1-12</sub> alkanyl, C<sub>7-20</sub> phenylalkyl, C<sub>11-20</sub> naphthylalkyl, C<sub>1-12</sub>, hydroxyalkyl, C<sub>2-12</sub> hydroxyalkenyl, C<sub>7-20</sub> hydroxyphenylalkyl, or C<sub>11-20</sub> hydroxynaphthylalkyl; or one and only one of R<sub>1</sub> and R<sub>2</sub> is COE<sub>1</sub> in which E<sub>1</sub> is C<sub>1-12</sub> alkyl, C<sub>2-12</sub> alkyl, C<sub>2-12</sub> alkenyl, C<sub>7-20</sub> phenylalkyl, C<sub>11-20</sub> naphthylalkyl, C<sub>1-12</sub> hydroxyalkyl, C<sub>2-12</sub> hydroxyalkenyl, C<sub>7-20</sub> hydroxyphenylalkyl, or C<sub>11-20</sub> hydroxynaphthylalkyl; and

10 R<sub>3</sub> is OH, NH<sub>2</sub>, C<sub>1-12</sub> alkoxy, or NH-Y-CH<sub>2</sub>-Z in which Y is a C<sub>1-12</sub> hydrocarbon moiety and Z is H, OH, CO<sub>2</sub>H or CONH<sub>2</sub>;

provided that (i) at least one of A<sub>5</sub>, A<sub>7</sub>, A<sub>8</sub>, A<sub>11</sub>, A<sub>15</sub>, A<sub>18</sub>, A<sub>22</sub>, A<sub>23</sub>, A<sub>24</sub>, A<sub>27</sub>, A<sub>28</sub>, A<sub>30</sub>, or A<sub>31</sub> is Cha, or at

15 least one of A<sub>3</sub>, A<sub>12</sub>, A<sub>16</sub>, A<sub>17</sub>, A<sub>18</sub>, A<sub>19</sub>, A<sub>22</sub>, A<sub>25</sub>, A<sub>26</sub>, A<sub>29</sub>, A<sub>30</sub>, or A<sub>34</sub> is Aib; or that (ii) at least one of A<sub>23</sub>, A<sub>24</sub>, A<sub>27</sub>, A<sub>28</sub>, or A<sub>31</sub> is Lys; or a pharmaceutically acceptable salt thereof. In one embodiment, at least one of A<sub>7</sub> and A<sub>11</sub> is Cha. In another embodiment, at least one

20 of A<sub>16</sub> or A<sub>19</sub> is Aib. Specific examples of peptides of the just-recited formula include, but are not limited to, [Cha<sup>7</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Cha<sup>11</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Cha<sup>7, 11</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Aib<sup>16</sup>, Tyr<sup>34</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Aib<sup>19</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Aib<sup>16, 19</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Cha<sup>7, 11</sup>, Aib<sup>16</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Cha<sup>7, 11</sup>, Aib<sup>19</sup>]hPTHrP(1-34)NH<sub>2</sub>;

25 [Cha<sup>22</sup>, Leu<sup>23, 28, 31</sup>, Glu<sup>25, 29</sup>, Lys<sup>26, 30</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Glu<sup>22, 25, 29</sup>, Leu<sup>23, 28, 31</sup>, Lys<sup>26, 27, 30</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Cha<sup>22, 23</sup>, Glu<sup>25, 29</sup>, Leu<sup>28, 31</sup>, Lys<sup>26, 30</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Glu<sup>22, 25</sup>, Leu<sup>23, 28, 31</sup>, Aib<sup>29</sup>, Lys<sup>26, 30</sup>]hPTHrP(1-34)NH<sub>2</sub>;

30 [Glu<sup>22, 25, 29</sup>, Lys<sup>23, 26, 30</sup>, Leu<sup>28, 31</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Glu<sup>22, 25, 29</sup>, Leu<sup>23, 28, 31</sup>, Lys<sup>26</sup>, Cha<sup>30</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Glu<sup>22, 25, 29</sup>, Leu<sup>23, 28, 31</sup>, Lys<sup>26</sup>, Aib<sup>30</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Glu<sup>22, 25, 29</sup>, Leu<sup>23, 31</sup>, Lys<sup>26, 28, 30</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Cha<sup>22, 23, 24, 27, 28, 31</sup>, Glu<sup>25, 29</sup>, Lys<sup>26, 30</sup>]hPTHrP(1-

35 34)NH<sub>2</sub>; [Glu<sup>22, 25, 29</sup>, Cha<sup>23, 24, 28, 31</sup>, Lys<sup>26, 27</sup>,

- 8 -

- <sup>30</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Glu<sup>22, 25, 29</sup>, Cha<sup>23, 24, 27, 31</sup>, Lys<sup>26, 28, 30</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Glu<sup>22, 25, 29</sup>, Lys<sup>23, 26, 30</sup>, Cha<sup>24, 27, 28, 31</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Cha<sup>22</sup>, Leu<sup>23, 28, 31</sup>, Glu<sup>25, 29</sup>, Lys<sup>26, 27, 30</sup>] hPTHrP(1-34)NH<sub>2</sub>; [Cha<sup>22</sup>, Leu<sup>23, 31</sup>, Glu<sup>25, 29</sup>, Lys<sup>26, 28, 30</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Cha<sup>22</sup>, Lys<sup>23, 26, 30</sup>, Glu<sup>25, 29</sup>, Leu<sup>28, 31</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Cha<sup>22</sup>, Leu<sup>23, 28, 31</sup>, Glu<sup>25, 29</sup>, Lys<sup>26, 30</sup>, Aib<sup>29</sup>] hPTHrP(1-34)NH<sub>2</sub>; [Cha<sup>22</sup>, Leu<sup>23, 28, 31</sup>, Glu<sup>25, 29</sup>, Lys<sup>26</sup>, Aib<sup>30</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Glu<sup>22, 25</sup>, Leu<sup>23, 28, 31</sup>, Lys<sup>26, 27, 30</sup>, Aib<sup>29</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Glu<sup>22, 25</sup>, Lys<sup>23, 26, 30</sup>, Leu<sup>28, 31</sup>, Aib<sup>29</sup>] hPTHrP(1-34)NH<sub>2</sub>; [Glu<sup>22, 25</sup>, Leu<sup>23, 31</sup>, Lys<sup>26, 28, 30</sup>, Aib<sup>29</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Cha<sup>7, 11</sup>, Glu<sup>22, 25, 29</sup>, Leu<sup>23, 28, 31</sup>, Lys<sup>26, 30</sup>] hPTHrP(1-34)NH<sub>2</sub>; [Cha<sup>7, 11, 22</sup>, Leu<sup>23, 28, 31</sup>, Glu<sup>25, 29</sup>, Lys<sup>26, 30</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Cha<sup>7, 11</sup>, Glu<sup>22, 25, 29</sup>, Leu<sup>23, 28, 31</sup>, Lys<sup>26, 27, 30</sup>] hPTHrP(1-34)NH<sub>2</sub>; [Cha<sup>7, 11, 22, 23</sup>, Glu<sup>25, 29</sup>, Leu<sup>28, 31</sup>, Lys<sup>26, 30</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Cha<sup>7, 11</sup>, Glu<sup>22, 25, 29</sup>, Lys<sup>23, 26, 30</sup>, Leu<sup>28, 31</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Cha<sup>7, 11</sup>, Glu<sup>22, 25, 29</sup>, Leu<sup>23, 31</sup>, Lys<sup>26, 28, 30</sup>] hPTHrP(1-34)NH<sub>2</sub>; [Cha<sup>7, 11</sup>, Glu<sup>22, 25</sup>, Leu<sup>23, 28, 31</sup>, Aib<sup>29</sup>, Lys<sup>26, 30</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Cha<sup>7, 11</sup>, Glu<sup>22, 25, 29</sup>, Leu<sup>23, 28, 31</sup>, Lys<sup>26</sup>, Aib<sup>30</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Cha<sup>15</sup>, Glu<sup>22, 25, 29</sup>, Leu<sup>23, 28, 31</sup>, Lys<sup>26, 30</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Cha<sup>15, 22</sup>, Leu<sup>23, 28, 31</sup>, Glu<sup>25, 29</sup>, Lys<sup>26, 30</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Cha<sup>15</sup>, Glu<sup>22, 25, 29</sup>, Leu<sup>23, 28, 31</sup>, Lys<sup>26, 27, 30</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Cha<sup>15, 22, 23</sup>, Glu<sup>25, 29</sup>, Leu<sup>28, 31</sup>, Lys<sup>26, 30</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Cha<sup>15</sup>, Glu<sup>22, 25</sup>, Leu<sup>23, 28, 31</sup>, Aib<sup>29</sup>, Lys<sup>26, 30</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Cha<sup>15</sup>, Glu<sup>22, 25, 29</sup>, Lys<sup>23, 26, 30</sup>, Leu<sup>28, 31</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Cha<sup>15</sup>, Glu<sup>22, 25, 29</sup>, Leu<sup>23, 28, 31</sup>, Lys<sup>26</sup>, Aib<sup>30</sup>] hPTHrP(1-34)NH<sub>2</sub>; [Cha<sup>15</sup>, Glu<sup>22, 28, 29</sup>, Leu<sup>23, 31</sup>, Lys<sup>26, 28</sup>, Aib<sup>30</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Cha<sup>15, 30</sup>, Glu<sup>22, 25, 29</sup>, Leu<sup>23, 28, 31</sup>, Lys<sup>26</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Cha<sup>7, 8, 22</sup>, Leu<sup>23, 28, 31</sup>, Glu<sup>25, 29</sup>, Lys<sup>26, 30</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Cha<sup>7, 8</sup>, Glu<sup>22, 25, 29</sup>, Leu<sup>23, 28, 31</sup>, Lys<sup>26, 27, 30</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Cha<sup>7, 8, 22, 23</sup>, Glu<sup>25, 29</sup>, Leu<sup>28, 31</sup>, Lys<sup>26, 30</sup>] hPTHrP (1-34)NH<sub>2</sub>; [Cha<sup>7, 8</sup>, Glu<sup>22, 25, 29</sup>, Leu<sup>23, 28, 31</sup>, Lys<sup>26, 30</sup>] hPTHrP(1-34)NH<sub>2</sub>; [Cha<sup>7, 8</sup>,

- 9 -

Glu<sup>22, 25</sup>, Leu<sup>23, 28, 31</sup>, Aib<sup>29</sup>, Lys<sup>26, 30</sup>]hPTHrP(1-34) NH<sub>2</sub>;  
 [Cha<sup>7, 8</sup>, Glu<sup>22, 25, 29</sup>, Lys<sup>23, 26, 30</sup>, Leu<sup>28, 31</sup>]hPTHrP(1-  
 34)NH<sub>2</sub>; [Cha<sup>7, 8</sup>, Glu<sup>22, 25, 29</sup>, Leu<sup>23, 28, 31</sup>, Lys<sup>26</sup>, Aib<sup>30</sup>]  
 hPTHrP(1-34)NH<sub>2</sub>; [Cha<sup>7, 8</sup>, Glu<sup>22, 25, 29</sup>, Leu<sup>23, 31</sup>, Lys<sup>26</sup>,  
 5 28, 30]hPTHrP(1-34)NH<sub>2</sub>; [Cha<sup>7, 8, 30</sup>, Glu<sup>22, 25, 29</sup>, Leu<sup>23, 28</sup>,  
 31, Lys<sup>26</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Ser<sup>1</sup>, Ile<sup>5</sup>, Cha<sup>7, 11, 22</sup>, Met<sup>8</sup>,  
 Asn<sup>10</sup>, His<sup>14</sup>, Leu<sup>23, 28, 31</sup>, Glu<sup>25, 29</sup>, Lys<sup>26, 30</sup>]hPTHrP(1-34)  
 NH<sub>2</sub>; [Ser<sup>1</sup>, Ile<sup>5</sup>, Cha<sup>7, 11</sup>, Met<sup>8</sup>, Asn<sup>10</sup>, His<sup>14</sup>, Glu<sup>22, 25</sup>,  
 29, Leu<sup>23, 28, 31</sup>, Lys<sup>26, 27, 30</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Ser<sup>1</sup>,  
 10 Ile<sup>5</sup>, Cha<sup>7, 11</sup>, Met<sup>8</sup>, Asn<sup>10</sup>, His<sup>14</sup>, Glu<sup>22, 25, 29</sup>, Leu<sup>23, 31</sup>,  
 Lys<sup>26, 28, 30</sup>]hPTHrP(1-34)NH<sub>2</sub>; Ser<sup>1</sup>, Ile<sup>5</sup>, Cha<sup>7, 11</sup>, Met<sup>8</sup>,  
 Asn<sup>10</sup>, His<sup>14</sup>, Glu<sup>22, 25, 29</sup>, Lys<sup>23, 26, 30</sup>, Leu<sup>28</sup>,  
 31]hPTHrP(1-34)NH<sub>2</sub>; [Ser<sup>1</sup>, Ile<sup>5</sup>, Cha<sup>7, 11</sup>, Met<sup>8</sup>, Asn<sup>10</sup>,  
 His<sup>14</sup>, Glu<sup>22, 25</sup>, Leu<sup>23, 28, 31</sup>, Aib<sup>29</sup>, Lys<sup>26, 30</sup>]hPTHrP(1-34)  
 15 NH<sub>2</sub>; [Ser<sup>1</sup>, Ile<sup>5</sup>, Cha<sup>7, 11</sup>, Met<sup>8</sup>, Asn<sup>10</sup>, His<sup>14</sup>, Glu<sup>22, 25</sup>,  
 29, Leu<sup>23, 28, 31</sup>, Lys<sup>26</sup>, Aib<sup>30</sup>] PTHrP(1-34)NH<sub>2</sub>; [Ser<sup>1</sup>,  
 Ile<sup>5</sup>, Cha<sup>7, 11, 22, 23</sup>, Met<sup>8</sup>, Asn<sup>10</sup>, His<sup>14</sup>, Glu<sup>25, 29</sup>, Leu<sup>28</sup>,  
 31, Lys<sup>26, 30</sup>]hPTHrP(1-34) NH<sub>2</sub>; [Ser<sup>1</sup>, Ile<sup>5</sup>, Cha<sup>7, 11, 15</sup>,  
 Met<sup>8</sup>, Asn<sup>10</sup>, His<sup>14</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Ser<sup>1</sup>, Ile<sup>5</sup>, Met<sup>8</sup>,  
 20 Asn<sup>10</sup>, Leu<sup>11</sup>, His<sup>14</sup>, Aib<sup>16</sup>]hPTHrP (1-34)NH<sub>2</sub>; [Ser<sup>1</sup>, Ile<sup>5</sup>,  
 Met<sup>8</sup>, Asn<sup>10</sup>, Leu<sup>11, 28, 31</sup>, His<sup>14</sup>, Cha<sup>22, 23</sup>, Glu<sup>25, 29</sup>,  
 Lys<sup>26, 30</sup>]hPTHrP (1-34)NH<sub>2</sub>; [Ser<sup>1</sup>, Ile<sup>5</sup>, Cha<sup>7, 11</sup>, Met<sup>8</sup>,  
 Asn<sup>10</sup>, His<sup>14</sup>, Glu<sup>22, 25, 29</sup>, Leu<sup>23, 28, 31</sup>, Lys<sup>26, 30</sup>]hPTHrP  
 (1-34)NH<sub>2</sub>; [Ser<sup>1</sup>, Ile<sup>5</sup>, Met<sup>8</sup>, Asn<sup>10</sup>, His<sup>14</sup>, Cha<sup>15</sup>, Glu<sup>22</sup>,  
 25, 29, Leu<sup>23, 28, 31</sup>, Lys<sup>26, 30</sup>]hPTHrP (1-34)NH<sub>2</sub>; [Ser<sup>1</sup>,  
 25 Ile<sup>5</sup>, Cha<sup>7, 8</sup>, Asn<sup>10</sup>, His<sup>14</sup>, Glu<sup>22, 25, 29</sup>, Leu<sup>23, 28, 31</sup>,  
 Lys<sup>26, 30</sup>]hPTHrP (1-34)NH<sub>2</sub>; [Glu<sup>22, 25, 29</sup>, Leu<sup>23, 28, 31</sup>,  
 Lys<sup>24, 26, 30</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Aib<sup>22</sup>, Leu<sup>23, 28, 31</sup>, Glu<sup>25</sup>,  
 29, Lys<sup>26, 30</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Glu<sup>22, 29</sup>, Leu<sup>23, 28, 31</sup>,  
 30 Aib<sup>25</sup>, Lys<sup>26, 30</sup>]hPTHrP(1-34)NH<sub>2</sub>; [Glu<sup>22, 25, 29</sup>, Leu<sup>23, 28</sup>,  
 31, Aib<sup>26</sup>, Lys<sup>30</sup>]hPTHrP (1-34)NH<sub>2</sub>; [Glu<sup>22, 25, 29</sup>, Leu<sup>23, 28</sup>,  
 Lys<sup>26, 30, 31</sup>] hPTHrP(1-34)NH<sub>2</sub>; [Ser<sup>1</sup>, Ile<sup>5</sup>, Met<sup>8</sup>, Asn<sup>10</sup>,  
 Leu<sup>11, 23, 28, 31</sup>, His<sup>14</sup>, Cha<sup>22</sup>, Glu<sup>25, 29</sup>, Lys<sup>26, 30</sup>]  
 hPTHrP(1-34)NH<sub>2</sub>; [Ser<sup>1</sup>, Ile<sup>5</sup>, Met<sup>8</sup>, Asn<sup>10</sup>, Leu<sup>11, 28, 31</sup>,  
 35 His<sup>14</sup>, Glu<sup>22, 25, 29</sup>, Lys<sup>23, 26, 30</sup>]PTHrP(1-34)NH<sub>2</sub>; [Ser<sup>1</sup>,

- 10 -

Ile<sup>5</sup>, Met<sup>8</sup>, Asn<sup>10</sup>, Leu<sup>11, 23, 28, 31</sup>, His<sup>14</sup>, Glu<sup>22, 25, 29</sup>, Lys<sup>26, 27, 30</sup>] hPTHrP(1-34)NH<sub>2</sub>; [Ser<sup>1</sup>, Ile<sup>5</sup>, Met<sup>8</sup>, Asn<sup>10</sup>, Leu<sup>11, 23, 31</sup>, His<sup>14</sup>, Glu<sup>22, 25, 29</sup>, Lys<sup>26, 28, 30</sup>] hPTHrP(1-34)NH<sub>2</sub>; [Ser<sup>1</sup>, Ile<sup>5</sup>, Met<sup>8</sup>, Asn<sup>10</sup>, Leu<sup>11, 23, 28, 31</sup>, His<sup>14</sup>, Glu<sup>22, 25</sup>, Aib<sup>29</sup>, Lys<sup>26, 30</sup>] hPTHrP(1-34)NH<sub>2</sub>; [Ser<sup>1</sup>, Ile<sup>5</sup>, Met<sup>8</sup>, Asn<sup>10</sup>, Leu<sup>11, 23, 28, 31</sup>, His<sup>14</sup>, Glu<sup>22, 25, 29</sup>, Lys<sup>26</sup>, Aib<sup>30</sup>] hPTHrP(1-34)NH<sub>2</sub>; or [Ser<sup>1</sup>, Ile<sup>5</sup>, Met<sup>8</sup>]hPTHrP(1-34)NH<sub>2</sub>.

With the exception of the N-terminal amino acid, all abbreviations (e.g. Ala or A<sub>1</sub>) of amino acids in this disclosure stand for the structure of -NH-CH(R)-CO-, wherein R is a side chain of an amino acid (e.g., CH<sub>3</sub> for Ala). For the N-terminal amino acid, the abbreviation stands for the structure of =N-CH(R)-CO-, wherein R is a side chain of an amino acid. β-Nal, Nle, Dap, Cha, Nva, Amp, Pal, and Aib are the abbreviations of the following α-amino acids: β-(2-naphthyl)alanine, norleucine, α,β-diaminopropionic acid, cyclohexylalanine, norvaline, 4-amino-phenylalanine, 3-pyridinylalanine, and α-aminoisobutyric acid, respectively. In the above formula, hydroxyalkyl, hydroxyphenyl-alkyl, and hydroxynaphthylalkyl may contain 1-4 hydroxy substituents. Also, COE<sub>1</sub> stands for -C=O·E<sub>1</sub>. Examples of -C=O·E<sub>1</sub> include, but are not limited to, acetyl and phenylpropionyl.

A peptide of this invention is also denoted herein by another format, e.g., [Cha<sup>7, 11</sup>]hPTH(1-34)NH<sub>2</sub>, with the substituted amino acids from the natural sequence placed between the second set of brackets (e.g., Cha<sup>7</sup> for Leu<sup>7</sup>, and Cha<sup>11</sup> for Leu<sup>11</sup> in hPTH). The abbreviation hPTH stands for human PTH, hPTHrP for human PTHrP, rPTH for rat PTH, and bPTH for bovine PTH. The numbers between the parentheses refer to the number of amino acids present in the peptide (e.g., hPTH(1-34) is amino acids 1 through 34 of the peptide sequence for human PTH). The

- 11 -

sequences for hPTH(1-34), hPTHrP(1-34), bPTH(1-34), and rPTH(1-34) are listed in Nissenson, et al., Receptor, 3:193 (1993). The designation "NH<sub>2</sub>" in PTH(1-34)NH<sub>2</sub> indicates that the C-terminus of the peptide is amidated. 5 PTH(1-34), on the other hand, has a free acid C-terminus.

Each of the peptides of the invention is capable of stimulating the growth of bone in a subject (i.e., a mammal such as a human patient). Thus, it is useful in the treatment of osteoporosis and bone fractures when 10 administered alone or concurrently with antiresorptive therapy, e.g., bisphosphonates and calcitonin.

The peptides of this invention can be provided in the form of pharmaceutically acceptable salts. Examples of such salts include, but are not limited to, those 15 formed with organic acids (e.g., acetic, lactic, maleic, citric, malic, ascorbic, succinic, benzoic, methanesulfonic, toluenesulfonic, or pamoic acid), inorganic acids (e.g., hydrochloric acid, sulfuric acid, or phosphoric acid), and polymeric acids (e.g., tannic 20 acid, carboxymethyl cellulose, polylactic, polyglycolic, or copolymers of polylactic-glycolic acids).

A therapeutically effective amount of a peptide of this invention and a pharmaceutically acceptable carrier substance (e.g., magnesium carbonate, lactose, or a 25 phospholipid with which the therapeutic compound can form a micelle) together form a therapeutic composition (e.g., a pill, tablet, capsule, or liquid) for administration (e.g., orally, intravenously, transdermally, pulmonarily, vaginally, subcutaneously, nasally, iontophoretically, or 30 by intratracheally) to a subject. The pill, tablet, or capsule that is to be administered orally can be coated with a substance for protecting the active composition from the gastric acid or intestinal enzymes in the stomach for a period of time sufficient to allow it to 35 pass undigested into the small intestine. The

- 12 -

therapeutic composition can also be in the form of a biodegradable or nonbiodegradable sustained release formulation for subcutaneous or intramuscular administration. See, e.g., U.S. Patents 3,773,919 and  
5 4,767,628 and PCT Application No. WO 94/15587.

Continuous administration can also be achieved using an implantable or external pump (e.g., INFUSAID™ pump). The administration can also be conducted intermittently, e.g., single daily injection, or continuously at a low  
10 dose, e.g., sustained release formulation.

The dose of a peptide of the present invention for treating the above-mentioned diseases or disorders varies depending upon the manner of administration, the age and the body weight of the subject, and the condition of the  
15 subject to be treated, and ultimately will be decided by the attending physician or veterinarian.

Also contemplated within the scope of this invention is a peptide covered by the above generic formula for use in treating diseases or disorders  
20 associated with deficiency in bone growth or the like, e.g., osteoporosis or fractures.

Other features and advantages of the present invention will be apparent from the detailed description and from the claims.

## 25        Detailed Description of the Invention

Based on the description herein, the present invention can be utilized to its fullest extent. The following specific examples are to be construed as merely illustrative, and not limitative of the remainder of the  
30 disclosure in any way whatsoever. Further, all publications cited herein are incorporated by reference.

### Structure

PTH(1-34) has been reported to have two amphophilic alpha helical domains. See, e.g., Barden, et

- 13 -

al., Biochem., 32:7126 (1992). The first  $\alpha$ -helix is formed between amino acid residues 4 through 13, while the second  $\alpha$ -helix is formed between amino acid residues 21 through 29. Some peptides of this invention contain  
5 the substitution of Cha for one or more residues within or near these two regions of PTH(1-34), e.g., Cha<sup>7</sup> and Cha<sup>11</sup> within the first  $\alpha$ -helix or Cha<sup>27</sup> and Cha<sup>28</sup> within the second  $\alpha$ -helix.

Also covered by this invention are variants of  
10 PTH(1-34) with the substitution of Aib for a residue adjacent to the  $\alpha$ -helixes, e.g., Aib<sup>16</sup>, Aib<sup>19</sup>, and Aib<sup>34</sup>; hArg<sup>27</sup> and Nle<sup>31</sup>, or the substitution of Dpa for the N-terminal residue.

#### Synthesis

15 The peptides of the invention can be prepared by standard solid phase synthesis. See, e.g., Stewart, J.M., et al., Solid Phase Synthesis (Pierce Chemical Co., 2d ed. 1984). The following is a description of how [Aib<sup>34</sup>]hPTH(1-34)NH<sub>2</sub> was prepared. Other peptides of the  
20 invention can be prepared in an analogous manner by a person of ordinary skill in the art.

The peptide was synthesized on an Applied Biosystems (Foster City, CA) model 430A peptide synthesizer which was modified to do accelerated Boc-  
25 chemistry solid phase peptide synthesis. See Schnoize, et al., Int. J. Peptide Protein Res., 90:180 (1992). 4-Methylbenz-hydrylamine (MBHA) resin (Peninsula, Belmont, CA) with the substitution of 0.93 mmol/g was used. The Boc amino acids (Bachem, CA, Torrance, CA; Nova Biochem.,  
30 LaJolla, CA) were used with the following side chain protection: Boc-Arg(Tos)-OH, Boc-Asp(OCHx1)-OH, Boc-Asn(Xan)-OH, Boc-Glu(OCHx1)-OH, Boc-His(DNP)-OH, Boc-Asn-GH, Boc-Val-OH, Boc-Leu-OH, Boc-Ser-OH, Boc-Gly-OH, Boc-Met-OH, Boc-Gln-OH, Boc-Ile-OH, Boc-Lys(2ClZ)-OH, Boc-

- 14 -

Ser(Bzl)-OH, and Boc-Trp(Fm)-OH. The synthesis was carried out on a 0.14 mmol scale. The Boc groups were removed by treatment with 100% TFA for 2 x 1 min. Boc amino acids (2.5 mmol) were pre-activated with HBTU (2.0 mmol) and DIEA (1.0 mL) in 4 mL of DMF and were coupled without prior neutralization of the peptide-resin TFA salt. Coupling times were 5 min except for the Boc-Aib-OH and the following residue, Boc-Asn(Xan)-OH, wherein the coupling times were 20 min.

At the end of the assembly of the peptide chain, the resin was treated with a solution of 20% mercaptoethanol/10% DIEA in DMF for 2 x 30 min. to remove the DNP group on the His side chain. The N-terminal Boc group was then removed by treatment with 100% TFA for 2 x 2 min. After neutralization of the peptide-resin with 10% DIEA in DMF (1 x 1 min.), the formyl group on the side chain of Trp was removed by treatment with a solution of 15% ethanolamine/15% water/70% DMF for 2 x 30 min. The partially-deprotected peptide-resin was washed with DMF and DCM and dried under reduced pressure. The final cleavage was done by stirring the peptide-resin in 10 mL of HF containing 1 mL of anisole at 0°C for 75 min. HF was removed by a flow of nitrogen. The residue was washed with ether (6 x 10 mL) and extracted with 4N HOAc (6 x 10 mL).

The peptide mixture in the aqueous extract was purified on a reversed-phase preparative high pressure liquid chromatography (HPLC) using a reversed phase Vydac™ C<sub>18</sub> column (Nest Group, Southborough, MA). The column was eluted with a linear gradient (10% to 45% of solution B over 130 min.) at a flow rate of 10 mL/min (Solution A = 0.1% aqueous TFA; Solution B = acetonitrile containing 0.1% of TFA). Fractions were collected and checked on analytical HPLC. Those containing pure product were combined and lyophilized to dryness. 62.3



- 15 -

mg of a white solid was obtained. Purity was >99% based on analytical HPLC analysis. Electro-spray mass spectrometer analysis gave the molecular weight at 4054.7 (in agreement with the calculated molecular weight of 4054.7).

The synthesis and purification of [Cha<sup>7,11</sup>]hPTH (1-34)NH<sub>2</sub> was carried out in the same manner as the above synthesis of [Aib<sup>34</sup>]hPTH(1-34)NH<sub>2</sub>. The protected amino acid Boc-Cha-OH was purchased from Bachem, CA. The purity of the final product was >98%, and the electron-spray mass spectrometer gave the molecular weight at 4197.0 (calculated molecular weight is 4196.9).

The full names for the abbreviations used above are as follows: Boc for t-butyloxycarbonyl, HF for hydrogen fluoride, Fm for formyl, Xan for xanthyl, Bzl for benzyl, Tos for tosyl, DNP for 2,4-dinitrophenyl, DMF for dimethylformamide, DCM for dichloromethane, HBTU for 2-(1H-Benzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate, DIEA for diisopropylethylamine, HOAc for acetic acid, TFA for trifluoroacetic acid, 2ClZ for 2-chlorobenzoyloxycarbonyl and OcHxl for O-cyclohexyl.

The substituents R<sub>1</sub> and R<sub>2</sub> of the above generic formula may be attached to the free amine of the N-terminal amino acid by standard methods known in the art. For example, alkyl groups, e.g., C<sub>1-12</sub> alkyl, may be attached using reductive alkylation. Hydroxyalkyl groups, e.g., C<sub>1-12</sub> hydroxyalkyl, may also be attached using reductive alkylation wherein the free hydroxy group is protected with a t-butyl ester. Acyl groups, e.g., COE<sub>1</sub>, may be attached by coupling the free acid, e.g., E<sub>1</sub>COOH, to the free amine of the N-terminal amino acid by mixing the completed resin with 3 molar equivalents of both the free acid and diisopropylcarbodiimide in methylene chloride for one hour and cycling the resulting resin through steps (a) to (f) in the above wash program. If the free acid contains a free hydroxy group, e.g., p-

- 16 -

hydroxyphenylpropionic acid, then the coupling should be performed with an additional 3 molar equivalents of HOBt.

Other peptides of this invention can be prepared in an analogous manner by a person of ordinary skill in the art.

### Functional Assays

#### A. Binding to PTH Receptor

The peptides of the invention were tested for their ability to bind to the PTH receptor present on SaOS-2 (human osteosarcoma cells). SaOS-2 cells (American Type Culture Collection, Rockville, MD; ATCC #HTB 85) were maintained in RPMI 1640 medium (Sigma, St. Louis, MO) supplemented with 10% fetal bovine serum (FBS) and 2 mM glutamine at 37°C in a humidified atmosphere of 5% CO<sub>2</sub> in air. The medium was changed every three or four days, and the cells were subcultured every week by trypsinization.

SaOS-2 cells were maintained for four days until they had reached confluence. The medium was replaced with 5% FBS in RPMI 1640 medium and incubated for 2 hrs at room temperature with  $10 \times 10^4$  cpm mono-<sup>125</sup>I-[Nle<sup>8,18</sup>, Tyr<sup>34</sup>(3-<sup>125</sup>I)] bPTH(1-34)NH<sub>2</sub> in the presence of a competing peptides of the invention at various concentrations between 10<sup>-11</sup>M to 10<sup>-4</sup> M. The cells were washed four times with ice-cold PBS and lysed with 0.1 M NaOH, and the radioactivity associated with the cells was counted in a scintillation counter. Synthesis of mono-<sup>125</sup>I-[Nle<sup>8,18</sup>, Tyr<sup>34</sup>(3-<sup>125</sup>I)] bPTH(1-34)NH<sub>2</sub> was carried out as described in Goldman, M.E., et al., Endocrinol., 123:1468 (1988).

The binding assay was conducted with various peptides of the invention, and the IC<sub>50</sub> value, (half maximal inhibition of binding of mono-<sup>125</sup>I-[Nle<sup>8,18</sup>, Tyr<sup>34</sup>(3-<sup>125</sup>I)]bPTH(1-34)NH<sub>2</sub>, for each peptide was calculated.

- 17 -

As shown in Table I, all of the tested peptides had a high binding affinity for the PTH receptor on the SaOS-2 cell.

#### B. Stimulation of Adenylate Cyclase Activity

5           The ability of the peptides of the invention to induce a biological response in SaOS-2 cells were measured. More specifically, any stimulation of the adenylate cyclase was determined by measuring the level of synthesis of cAMP (adenosine 3',5'-monophosphate) as  
10 described previously in Rodan, et al., J. Clin. Invest. 72: 1511 (1983) and Goldman, et al., Endocrinol., 123:1468 (1988). Confluent SAOS-2 cells in 24 wells plates were incubated with 0.5  $\mu$ Ci [ $^3$ H]adenine (26.9 Ci/mmol, New England Nuclear, Boston, MA) in fresh medium  
15 at 37°C for 2 hrs, and washed twice with Hank's balanced salt solution (Gibco, Gaithersburg, MD). The cells were treated with 1 mM IBMX [isobutylmethyl-xanthine, Sigma, St. Louis, MO] in fresh medium for 15 min, and the peptides of the invention were added to the medium to  
20 incubate for 5 min. The reaction was stopped by the addition of 1.2 M trichloroacetic acid (TCA) (Sigma, St. Louis, MO) followed by sample neutralization with 4 N KOH. cAMP was isolated by the two-column chromatographic method (Salmon, et al., 1974, Anal. Biochem. 58, 541).  
25 The radioactivity was counted in a scintillation counter (Liquid Scintillation Counter 2200CA, PACKARD, Downers Grove, IL).

          The respective EC<sub>50</sub> values (half maximal stimulation of adenylate cyclase) for the tested peptides  
30 were calculated and shown in Table I. All tested peptides were found to be potent stimulators of adenylate cyclase activity, which is a biochemical pathway indicative as a proximal signal for osteoblast proliferation (e.g., bone growth).

TABLE I

	PEPTIDE	K <sub>d</sub> (μM)	EC <sub>50</sub> (nM)
5	[Cha <sup>7, 11</sup> ]hPTH(1-34)NH <sub>2</sub>	0.01	0.6
	[Cha <sup>23</sup> ]hPTH(1-34)NH <sub>2</sub>	0.2	20
	[Cha <sup>24</sup> ]hPTH(1-34)NH <sub>2</sub>	0.1	10
	[Nle <sup>8, 18</sup> , Cha <sup>27</sup> ]hPTH(1-34)NH <sub>2</sub>	0.05	2
	[Cha <sup>28</sup> ]hPTH(1-34)NH <sub>2</sub>	0.05	2.5
10	[Cha <sup>31</sup> ]hPTH(1-34)NH <sub>2</sub>	0.03	4
	[Aib <sup>16</sup> ]hPTH(1-34)NH <sub>2</sub>	0.004	0.7
	[Aib <sup>19</sup> ]hPTH(1-34)NH <sub>2</sub>	0.005	0.6
	[Aib <sup>34</sup> ]hPTH(1-34)NH <sub>2</sub>	0.007	3
	[Nle <sup>31</sup> ]hPTH(1-34)NH <sub>2</sub>	0.004	0.7
15	[hArg <sup>27</sup> ]hPTH(1-34)NH <sub>2</sub>	0.007	1
	[Dap, Nle <sup>8, 18</sup> , Tyr <sup>34</sup> ]hPTH(1-34)NH <sub>2</sub>	0.150	10
	[Cha <sup>24, 28, 31</sup> , Lys <sup>30</sup> ]hPTH(1-34)NH <sub>2</sub>	0.5	7
	[Cha <sup>7, 11</sup> , Nle <sup>8, 18</sup> , Tyr <sup>34</sup> ]hPTH(1-34)NH <sub>2</sub>	0.006	0.6
	[Cha <sup>7, 11</sup> , Nle <sup>8, 18</sup> , Aib <sup>16, 19</sup> , Tyr <sup>34</sup> ]hPTH(1-34)NH <sub>2</sub>	0.005	1.5
20	[Cha <sup>7, 11</sup> , Nle <sup>8, 18, 31</sup> , Aib <sup>16, 19</sup> , Tyr <sup>34</sup> ]hPTH(1-34)NH <sub>2</sub>	0.04	4
	[Cha <sup>11</sup> ]hPTH(1-34)NH <sub>2</sub>	0.005	2
	[Cha <sup>28, 31</sup> ]hPTH(1-34)NH <sub>2</sub>	0.06	7
	[Cha <sup>7, 11</sup> , Nle <sup>8, 18</sup> , Aib <sup>34</sup> ]hPTH(1-34)NH <sub>2</sub>	0.03	1.5
	[Cha <sup>15</sup> ]hPTH(1-34)NH <sub>2</sub>	0.005	1.3
25	[Cha <sup>7, 11</sup> , Aib <sup>19</sup> ]hPTH(1-34)NH <sub>2</sub>	0.007	0.5
	[Cha <sup>7, 11</sup> , Aib <sup>16</sup> ]hPTH(1-34)NH <sub>2</sub>	0.004	1.1
	[Aib <sup>16, 19</sup> ]hPTH(1-34)NH <sub>2</sub>	0.004	0.6
	[Aib <sup>12</sup> ]hPTH(1-34)NH <sub>2</sub>	0.005	2
	[Aib <sup>3</sup> ]hPTH(1-34)NH <sub>2</sub>	0.004	1.1
30	[Cha <sup>7, 11</sup> , Aib <sup>19</sup> , Lys <sup>30</sup> ]hPTH(1-34)NH <sub>2</sub>	0.004	2
	[Cha <sup>7</sup> ]hPTH(1-34)NH <sub>2</sub>	0.02	2.3
	[Cha <sup>24, 28, 31</sup> ]hPTH(1-34)NH <sub>2</sub>	1.0	30
	[Aib <sup>17</sup> ]hPTH(1-34)	0.05	3
	[Cha <sup>7, 11, 15</sup> ]hPTH(1-34)	0.01	1.4

- 19 -


Other Embodiments

It is to be understood that while the invention has been described in conjunction with the detailed description thereof, that the foregoing description is  
5 intended to illustrate and not limit the scope of the invention, which is defined by the scope of the appended claims. Other aspects, advantages, and modifications are within the claims.

- 20 -

What is claimed is:

1. A peptide of the formula:



10 wherein

$A_1$  is Ser, Ala, or Dap;

$A_3$  is Ser, Thr, or Aib;

$A_5$  is Leu, Nle, Ile, Cha,  $\beta$ -Nal, Trp, Pal, Phe or p-X-Phe, in which X is OH, a halogen, or  $\text{CH}_3$ ;

15  $A_7$  is Leu, Nle, Ile, Cha,  $\beta$ -Nal, Trp, Pal, Phe, or p-X-Phe in which X is OH, a halogen, or  $\text{CH}_3$ ;

$A_8$  is Met, Nva, Leu, Val, Ile, Cha, or Nle;

$A_{11}$  is Leu, Nle, Ile, Cha,  $\beta$ -Nal, Trp, Pal, Phe or p-X-Phe in which X is OH, a halogen, or  $\text{CH}_3$ ;

20  $A_{12}$  is Gly or Aib;

$A_{15}$  is Leu, Nle, Ile, Cha,  $\beta$ -Nal, Trp, Pal, Phe, or p-X-Phe in which X is OH, a halogen, or  $\text{CH}_3$ ;

$A_{16}$  is Ser, Asn, Ala, or Aib;

$A_{17}$  is Ser, Thr, or Aib;

25  $A_{18}$  is Met, Nva, Leu, Val, Ile, Nle, Cha, or Aib;

$A_{19}$  is Glu or Aib;

$A_{21}$  is Val, Cha, or Met;

$A_{23}$  is Trp or Cha;

$A_{24}$  is Leu or Cha;

30  $A_{27}$  is Lys, Aib, Leu, hArg, Gln, or Cha;

$A_{28}$  is Leu or Cha;

$A_{30}$  is Asp or Lys;

$A_{31}$  is Val, Nle, Cha, or deleted;

$A_{32}$  is His or deleted;

- 21 -

- A<sub>33</sub> is Asn or deleted;  
A<sub>34</sub> is Phe, Tyr, Amp, Aib, or deleted;  
each of R<sub>1</sub> and R<sub>2</sub> is, independently, H, C<sub>1-12</sub> alkyl, C<sub>2-12</sub> alkenyl, C<sub>7-20</sub> phenylalkyl, C<sub>11-20</sub> naphthylalkyl, C<sub>1-12</sub> hydroxyalkyl, C<sub>2-12</sub> hydroxyalkenyl, C<sub>7-20</sub> hydroxyphenylalkyl, or C<sub>11-20</sub> hydroxynaphthylalkyl; or one and only one of R<sub>1</sub> and R<sub>2</sub> is COE<sub>1</sub> in which E<sub>1</sub> is C<sub>1-12</sub> alkyl, C<sub>2-12</sub> alkenyl, C<sub>7-20</sub> phenylalkyl, C<sub>11-20</sub> naphthylalkyl, C<sub>1-12</sub> hydroxyalkyl, C<sub>2-12</sub> hydroxyalkenyl, C<sub>7-20</sub> hydroxy-phenylalkyl, or C<sub>11-20</sub> hydroxynaphthylalkyl; and R<sub>3</sub> is OH, NH<sub>2</sub>, C<sub>1-12</sub> alkoxy, or NH-Y-CH<sub>2</sub>-Z in which Y is a C<sub>1-12</sub> hydrocarbon moiety and Z is H, OH, CO<sub>2</sub>H, or CONH<sub>2</sub>;
- provided that at least one of A<sub>5</sub>, A<sub>7</sub>, A<sub>8</sub>, A<sub>11</sub>, A<sub>15</sub>, A<sub>18</sub>, A<sub>21</sub>, A<sub>23</sub>, A<sub>24</sub>, A<sub>27</sub>, A<sub>28</sub>, and A<sub>31</sub> is Cha, or at least one of A<sub>3</sub>, A<sub>12</sub>, A<sub>16</sub>, A<sub>17</sub>, A<sub>18</sub>, A<sub>19</sub>, and A<sub>34</sub> is Aib; or a pharmaceutically acceptable salt thereof.

2. A peptide of claim 1, wherein at least one of A<sub>7</sub>, A<sub>11</sub>, A<sub>15</sub>, A<sub>23</sub>, A<sub>24</sub>, A<sub>27</sub>, A<sub>28</sub>, and A<sub>31</sub> is Cha; or a pharmaceutically acceptable salt thereof.

3. A peptide of claim 2, wherein

A<sub>3</sub> is Ser;  
A<sub>5</sub> is Ile;  
A<sub>7</sub> is Leu or Cha;  
A<sub>8</sub> is Met, Nva, Leu, Val, Ile, or Nle;  
A<sub>11</sub> is Leu or Cha;  
A<sub>12</sub> is Gly;  
A<sub>15</sub> is Leu or Cha;  
A<sub>16</sub> is Asn or Aib;  
A<sub>17</sub> is Ser;  
A<sub>18</sub> is Met or Nle;  
A<sub>21</sub> is Val;  
A<sub>27</sub> is Lys, hArg, or Cha;

- 22 -

A<sub>32</sub> is His;  
A<sub>31</sub> is Val, Nle, or Cha;  
A<sub>33</sub> is Asn;  
A<sub>34</sub> is Phe, Tyr, Amp, or Aib;  
5 R<sub>1</sub> is H;  
R<sub>2</sub> is H; and  
R<sub>3</sub> is NH<sub>2</sub>;  
or a pharmaceutically acceptable salt thereof.

4. A peptide of claim 3, wherein at least one of  
10 A<sub>7</sub> and A<sub>11</sub> is Cha; or a pharmaceutically acceptable salt thereof.

5. A peptide of claim 4, wherein said peptide is  
[Cha<sup>7, 11</sup>]hPTH(1-34)NH<sub>2</sub>, [Cha<sup>7, 11</sup>, Nle<sup>8, 18</sup>, Tyr<sup>34</sup>]hPTH(1-  
34)NH<sub>2</sub>; [Cha<sup>11</sup>]hPTH(1-34)NH<sub>2</sub>; [Cha<sup>7, 11, 15</sup>]hPTH(1-34)NH<sub>2</sub>; or  
15 [Cha<sup>7</sup>]hPTH(1-34)NH<sub>2</sub>; or a pharmaceutically acceptable salt thereof.

6. A peptide of claim 3, wherein at least one of  
A<sub>15</sub>, A<sub>23</sub>, A<sub>24</sub>, A<sub>27</sub>, A<sub>28</sub>, and A<sub>31</sub> is Cha; or a  
pharmaceutically acceptable salt thereof.

20 7. A peptide of claim 6, wherein said peptide is  
[Cha<sup>23</sup>]hPTH(1-34)NH<sub>2</sub>, [Cha<sup>24</sup>]hPTH(1-34)NH<sub>2</sub>, [Nle<sup>8, 18</sup>,  
Cha<sup>27</sup>]hPTH(1-34)NH<sub>2</sub>, [Cha<sup>28</sup>]hPTH(1-34)NH<sub>2</sub>, [Cha<sup>31</sup>]hPTH(1-  
34)NH<sub>2</sub>, [Cha<sup>24, 28, 31</sup>]hPTH(1-34)NH<sub>2</sub>; [Cha<sup>24, 28, 31</sup>,  
Lys<sup>30</sup>]hPTH(1-34)NH<sub>2</sub>; [Cha<sup>28, 31</sup>]hPTH(1-34)NH<sub>2</sub>; or  
25 [Cha<sup>15</sup>]hPTH(1-34)NH<sub>2</sub>; or a pharmaceutically acceptable  
salt thereof.

8. A peptide of claim 1, wherein at least one of  
A<sub>3</sub>, A<sub>12</sub>, A<sub>16</sub>, A<sub>17</sub>, A<sub>18</sub>, A<sub>19</sub>, and A<sub>34</sub> is Aib; or a  
pharmaceutically acceptable salt thereof.



- 23 -

9. A peptide of claim 8, wherein  
A<sub>3</sub> is Ser or Aib;  
A<sub>5</sub> is Ile;  
A<sub>7</sub> is Leu or Cha;  
5 A<sub>8</sub> is Met, Nva, Leu, Val, Ile, or Nle;  
A<sub>11</sub> is Leu or Cha;  
  
A<sub>15</sub> is Leu or Cha;  
A<sub>16</sub> is Asn or Aib;  
A<sub>18</sub> is Met, Aib, or Nle;  
10 A<sub>21</sub> is Val;  
A<sub>27</sub> is Lys, Aib, Leu, hArg, or Cha;  
A<sub>31</sub> is Val, Nle, or Cha;  
A<sub>32</sub> is His;  
A<sub>33</sub> is Asn;  
15 A<sub>34</sub> is Phe, Tyr, Amp, or Aib;  
R<sub>1</sub> is H;  
R<sub>2</sub> is H; and  
R<sub>3</sub> is NH<sub>2</sub>;

or a pharmaceutically acceptable salt thereof.

- 20 10. A peptide of claim 9, wherein at least one of  
A<sub>3</sub>, A<sub>12</sub>, A<sub>16</sub>, A<sub>17</sub>, A<sub>19</sub>, and A<sub>34</sub> is Aib; or a  
pharmaceutically acceptable salt thereof.

11. A peptide of claim 10, wherein said peptide  
is [Aib<sup>16</sup>]hPTH(1-34)NH<sub>2</sub>, [Aib<sup>19</sup>]hPTH(1-34)NH<sub>2</sub>,  
25 [Aib<sup>34</sup>]hPTH(1-34)NH<sub>2</sub>; [Aib<sup>16, 19</sup>]hPTH(1-34)NH<sub>2</sub>;  
[Aib<sup>3</sup>]hPTH(1-34)NH<sub>2</sub>; [Aib<sup>17</sup>]hPTH(1-34)NH<sub>2</sub>; or  
[Aib<sup>12</sup>]hPTH(1-34)NH<sub>2</sub>; or a pharmaceutically acceptable  
salt thereof.

12. A peptide of claim 1 wherein at least one of  
30 A<sub>7</sub>, A<sub>11</sub>, A<sub>15</sub>, A<sub>23</sub>, A<sub>24</sub>, A<sub>27</sub>, A<sub>28</sub>, and A<sub>31</sub> is Cha and at

- 24 -

least one of A<sub>3</sub>, A<sub>12</sub>, A<sub>16</sub>, A<sub>17</sub>, A<sub>18</sub>, A<sub>19</sub>, and A<sub>34</sub> is Aib;  
or a pharmaceutically acceptable salt thereof.

13. A peptide of claim 12, wherein
- 5 A<sub>3</sub> is Ser or Aib;  
A<sub>5</sub> is Ile;  
A<sub>7</sub> is Leu or Cha;  
A<sub>8</sub> is Met, Nva, Leu, Val, Ile, or Nle;  
A<sub>11</sub> is Leu or Cha;  
A<sub>15</sub> is Leu or Cha;  
10 A<sub>16</sub> is Asn or Aib;  
A<sub>18</sub> is Met, Aib, or Nle;  
A<sub>21</sub> is Val;  
A<sub>27</sub> is Lys, Aib, Leu, hArg, or Cha;  
A<sub>31</sub> is Val, Nle, or Cha;  
15 A<sub>32</sub> is His;  
A<sub>33</sub> is Asn;  
A<sub>34</sub> is Phe, Tye, Amp, or Aib;  
R<sub>1</sub> is H;  
R<sub>2</sub> is H; and  
20 R<sub>3</sub> is NH<sub>2</sub>;  
or a pharmaceutically acceptable salt thereof.

14. A peptide of claim 13, wherein at least one  
of A<sub>7</sub> and A<sub>11</sub> is Cha and at least one of A<sub>16</sub>, A<sub>19</sub>, and A<sub>34</sub>  
is Aib; or a pharmaceutically acceptable salt thereof.


- 25 15. A peptide of claim 14, wherein said peptide  
is [Cha<sup>7, 11</sup>, Nle<sup>8, 18</sup>, Aib<sup>16, 19</sup>, Tyr<sup>34</sup>]hPTH(1-34)NH<sub>2</sub>,  
[Cha<sup>7, 11</sup>, Nle<sup>8, 18, 31</sup>, Aib<sup>16, 19</sup>, Tyr<sup>34</sup>]hPTH(1-34)NH<sub>2</sub>;  
[Cha<sup>7, 11</sup>, Aib<sup>19</sup>]hPTH(1-34)NH<sub>2</sub>; [Cha<sup>7, 11</sup>, Aib<sup>16</sup>]hPTH(1-  
34)NH<sub>2</sub>; [Cha<sup>7, 11</sup>, Nle<sup>8, 18</sup>, Aib<sup>34</sup>]hPTH(1-34)NH<sub>2</sub>; or [Cha<sup>7,</sup>  
30 <sup>11</sup>, Aib<sup>19</sup>, Lys<sup>30</sup>]hPTH(1-34)NH<sub>2</sub>; or a pharmaceutically  
acceptable salt thereof.

- 25 -

16. A peptide of claim 13, wherein at least one of A<sub>24</sub>, A<sub>28</sub>, and A<sub>31</sub> is Cha and at least one of A<sub>16</sub> and A<sub>17</sub> is Aib; or a pharmaceutically acceptable salt thereof.

17. A peptide of claim 16, wherein said peptide is [Cha<sup>28</sup>, Nle<sup>8, 18</sup>, Aib<sup>16, 19</sup>, Tyr<sup>34</sup>]hPTH(1-34)NH<sub>2</sub>, or [Cha<sup>28</sup>, Aib<sup>16, 19</sup>] PTH(1-34)NH<sub>2</sub>; or a pharmaceutically acceptable salt thereof.

18. A peptide of the formula:



wherein

A<sub>3</sub> is Ser, Thr, or Aib;

A<sub>5</sub> is Leu, Nle, Ile, Cha, β-Nal, Trp, Pal, Phe or p-X-Phe, in which X is OH, a halogen, or CH<sub>3</sub>;

20 A<sub>7</sub> is Leu, Ile, Nle, Cha, β-Nal, Trp, Pal, Phe, or p-X-Phe in which X is H, OH, a halogen, or CH<sub>3</sub>;

A<sub>8</sub> is Met, Nva, Leu, Val, Ile, Cha, or Nle;

A<sub>11</sub> is Leu, Nle, Ile, Cha, β-Nal, Trp, Pal, Phe or p-X-Phe in which X is OH, a halogen, or CH<sub>3</sub>;

25 A<sub>12</sub> is Gly or Aib;

A<sub>15</sub> is Leu, Nle, Ile, Cha, β-Nal, Trp, Pal, Phe, or p-X-Phe in which X is OH, a halogen, or CH<sub>3</sub>;

A<sub>16</sub> is Ser, Asn, Ala, or Aib;

A<sub>17</sub> is Ser, Thr, or Aib;

30 A<sub>18</sub> is Met, Nva, Leu, Val, Ile, Nle, Cha, or Aib;

A<sub>19</sub> is Glu or Aib;

A<sub>21</sub> is Val, Cha, or Met;

A<sub>23</sub> is Trp or Cha;

- 26 -

- A<sub>24</sub> is Leu or Cha;  
A<sub>27</sub> is Lys, Aib, Leu, hArg, Gln, or Cha;  
A<sub>28</sub> is Leu or Cha;  
A<sub>30</sub> is Asp or Lys;  
5 A<sub>31</sub> is Val, Nle, Cha, or deleted;  
A<sub>32</sub> is His or deleted;  
A<sub>33</sub> is Asn or deleted;  
A<sub>34</sub> is Phe, Tyr, Amp, Aib, or deleted;  
each of R<sub>1</sub> and R<sub>2</sub> is, independently, H, C<sub>1-12</sub>  
10 alkyl, C<sub>2-12</sub> alkenyl, C<sub>7-20</sub> phenylalkyl, C<sub>11-20</sub>  
naphthylalkyl, C<sub>1-12</sub> hydroxyalkyl, C<sub>2-12</sub> hydroxyalkenyl, C<sub>7-20</sub>  
hydroxyphenylalkyl, or C<sub>11-20</sub> hydroxynaphthylalkyl; or  
one and only one of R<sub>1</sub> and R<sub>2</sub> is COE<sub>1</sub> in which E<sub>1</sub> is C<sub>1-12</sub>  
alkyl, C<sub>2-12</sub> alkenyl, C<sub>7-20</sub> phenylalkyl, C<sub>11-20</sub>  
15 naphthylalkyl, C<sub>1-12</sub> hydroxyalkyl, C<sub>2-12</sub> hydroxyalkenyl, C<sub>7-20</sub>  
hydroxy-phenylalkyl, or C<sub>11-20</sub> hydroxynaphthylalkyl;  
R<sub>3</sub> is OH, NH<sub>2</sub>, C<sub>1-12</sub> alkoxy, or NH-Y-CH<sub>2</sub>-Z in which  
Y is a C<sub>1-12</sub> hydrocarbon moiety and Z is H, OH, CO<sub>2</sub>H, or  
CONH<sub>2</sub>;  
20 provided that at least A<sub>1</sub> is Dap, A<sub>7</sub> is  $\beta$ -Nal,  
Trp, Pal, Phe, or p-X-Phe; A<sub>15</sub> is  $\beta$ -Nal, Trp, Pal, Phe, or  
p-X-Phe, A<sub>27</sub> is hArg, or A<sub>31</sub> is Nle; or a pharmaceutically  
acceptable salt thereof.

19. A peptide of claim 18, wherein  
25 A<sub>1</sub> is Ser, Gly, or Dap;  
A<sub>3</sub> is Ser or Aib;  
A<sub>8</sub> is Met, Nva, Leu, Val, Ile, or Nle;  
A<sub>16</sub> is Asn or Aib;  
A<sub>18</sub> is Met, Aib, or Nle;  
30 A<sub>21</sub> is Val;  
A<sub>27</sub> is Lys, Aib, Leu, hArg, or Cha;  
A<sub>31</sub> is Val, Nle, or Cha;  
A<sub>32</sub> is His;  
A<sub>33</sub> is Asn;

- 27 -

A<sub>34</sub> is Phe, Tyr, Amp, or Aib;

R<sub>1</sub> is H;


R<sub>2</sub> is H; and

R<sub>3</sub> is NH<sub>2</sub>;

5 or a pharmaceutically acceptable salt thereof.

20. A peptide of claim 19, wherein said peptide is [Nle<sup>31</sup>]hPTH(1-34)NH<sub>2</sub>, [hArg<sup>27</sup>]hPTH(1-34)NH<sub>2</sub>, or [Dap<sup>1</sup>, Nle<sup>8, 18</sup>, Tyr<sup>34</sup>]hPTH(1-34)NH<sub>2</sub>; or a pharmaceutically acceptable salt thereof.

10 21. A peptide of the formula:



wherein

20 A<sub>1</sub> is Ala, Ser, or Dap;

A<sub>3</sub> is Ser or Aib;

A<sub>5</sub> is His, Ile, or Cha;

A<sub>7</sub> is Leu, Cha, Nle, β-Nal, Trp, Pal, Phe, or p-X-Phe in which X is OH, a halogen, or CH<sub>3</sub>;

A<sub>8</sub> is Leu, Met, or Cha;

25 A<sub>10</sub> is Asp or Asn;

A<sub>11</sub> is Lys, Leu, Cha, Phe, or β-Nal;

A<sub>12</sub> is Gly or Aib;

A<sub>14</sub> is Ser or His;

A<sub>15</sub> is Ile, or Cha;

30 A<sub>16</sub> is Gln or Aib;

A<sub>17</sub> is Asp or Aib;

A<sub>18</sub> is Leu, Aib, or Cha;

- 28 -

- $A_{19}$  is Arg or Aib;  
 $A_{22}$  is Phe, Glu, Aib, or Cha;  
 $A_{23}$  is Phe, Leu, Lys, or Cha;  
 $A_{24}$  is Leu, Lys, or Cha;  
5  $A_{25}$  is His, Aib, or Glu;  
 $A_{26}$  is His, Aib, or Lys;  
 $A_{27}$  is Leu, Lys, or Cha;  
 $A_{28}$  is Ile, Leu, Lys, or Cha;  
 $A_{29}$  is Ala, Glu, or Aib;  
10  $A_{30}$  is Glu, Cha, Aib, or Lys;  
 $A_{31}$  is Ile, Leu, Cha, Lys, or deleted;  
 $A_{32}$  is His or deleted;  
 $A_{33}$  is Thr or deleted;  
 $A_{34}$  is Ala or deleted;  
15 each of  $R_1$  and  $R_2$  is, independently, H,  $C_{1-12}$  alkanyl,  $C_{7-20}$  phenylalkyl,  $C_{11-20}$  naphthylalkyl,  $C_{1-12}$  hydroxyalkyl,  $C_{2-12}$  hydroxyalkenyl,  $C_{7-20}$  hydroxyphenylalkyl, or  $C_{11-20}$  hydroxynaphthylalkyl; or one and only one of  $R_1$  and  $R_2$  is  $COE_1$  in which  $E_1$  is  $C_{1-12}$   
20 alkyl,  $C_{2-12}$  alkyl,  $C_{2-12}$  alkenyl,  $C_{7-20}$  phenylalkyl,  $C_{11-20}$  naphthylalkyl,  $C_{1-12}$  hydroxyalkyl,  $C_{2-12}$  hydroxyalkenyl,  $C_{7-20}$  hydroxyphenylalkyl, or  $C_{11-20}$  hydroxynaphthylalkyl; and  
 $R_3$  is OH,  $NH_2$ ,  $C_{1-12}$  alkoxy, or  $NH-Y-CH_2-Z$  in which  
Y is a  $C_{1-12}$  hydrocarbon moiety and Z is H, OH,  $CO_2H$  or  
25  $CONH_2$ ;  
provided that at least one of  $A_5$ ,  $A_7$ ,  $A_8$ ,  $A_{11}$ ,  $A_{15}$ ,  
 $A_{18}$ ,  $A_{22}$ ,  $A_{23}$ ,  $A_{24}$ ,  $A_{27}$ ,  $A_{28}$ ,  $A_{30}$ , or  $A_{31}$  is Cha, or at  
least one of  $A_3$ ,  $A_{12}$ ,  $A_{16}$ ,  $A_{17}$ ,  $A_{18}$ ,  $A_{19}$ ,  $A_{22}$ ,  $A_{25}$ ,  $A_{26}$ ,  
 $A_{29}$ ,  $A_{30}$ , or  $A_{34}$  is Aib; or a pharmaceutically acceptable  
30 salt thereof.

22. A peptide of claim 21, wherein at  $A_{22}$  is Phe or Cha;  $A_{23}$  is Phe or Cha;  $A_{25}$  is His;  $A_{26}$  is His;  $A_{27}$  is Leu or Cha;  $A_{28}$  is Ile or Cha;  $A_{29}$  is Ala;  $A_{30}$  is Glu or

- 29 -

Lys; A<sub>31</sub> is Ile or Cha; A<sub>32</sub> is His; A<sub>33</sub> is Thr; and A<sub>34</sub> is Ala; or a pharmaceutically acceptable salt thereof.

23. A peptide of claim 22, wherein at least one of A<sub>7</sub> and A<sub>11</sub> is Cha; or a pharmaceutically acceptable salt thereof.


24. A peptide of claim 22, wherein at least one of A<sub>16</sub> or A<sub>19</sub> is Aib; or a pharmaceutically acceptable salt thereof.

25. A peptide of claim 21, wherein A<sub>22</sub> is Glu, Aib, or Cha; A<sub>23</sub> is Leu, Lys, or Cha; A<sub>25</sub> is Aib or Glu; A<sub>26</sub> is Aib or Lys; A<sub>28</sub> is Leu, Lys, or Cha; A<sub>29</sub> is Glu or Aib; A<sub>30</sub> is Cha, Aib, or Lys; A<sub>31</sub> is Leu, Cha, or Lys; A<sub>32</sub> is His; A<sub>33</sub> is Thr; and A<sub>34</sub> is Ala; or a pharmaceutically acceptable salt thereof.

26. A peptide of claim 25, wherein at least one of A<sub>7</sub> and A<sub>11</sub> is Cha; or a pharmaceutically acceptable salt thereof.

27. A peptide of claim 25, wherein at least one of A<sub>16</sub> or A<sub>19</sub> is Aib; or a pharmaceutically acceptable salt thereof.

28. A peptide of the formula:



wherein

- 30 -

- $A_1$  is Ala, Ser, or Dap;  
 $A_3$  is Ser or Aib;  
 $A_5$  is His, Ile, or Cha;  
 $A_7$  is Leu, Cha, Nle,  $\beta$ -Nal, Trp, Pal, Phe, or  
5 p-X-Phe in which X is OH, a halogen, or  $CH_3$ ;  
 $A_8$  is Leu, Met, or Cha;  
 $A_{10}$  is Asp or Asn;  
 $A_{11}$  is Lys, Leu, Cha, Phe, or  $\beta$ -Nal;  
 $A_{12}$  is Gly or Aib;  
10  $A_{14}$  is Ser or His;  
 $A_{15}$  is Ile, or Cha;  
 $A_{16}$  is Gln or Aib;  
 $A_{17}$  is Asp or Aib;  
 $A_{18}$  is Leu, Aib, or Cha;  
15  $A_{19}$  is Arg or Aib;  
 $A_{22}$  is Phe, Glu, Aib, or Cha;  
 $A_{23}$  is Phe, Leu, Lys, or Cha;  
 $A_{24}$  is Leu, Lys, or Cha;  
 $A_{25}$  is His, Aib, or Glu;  
20  $A_{26}$  is His, Aib, or Lys;  
 $A_{27}$  is Leu, Lys, or Cha;  
 $A_{28}$  is Ile, Leu, Lys, or Cha;  
 $A_{29}$  is Ala, Glu, or Aib;  
 $A_{30}$  is Glu, Cha, Aib, or Lys;  
25  $A_{31}$  is Ile, Leu, Cha, Lys, or deleted;  
 $A_{32}$  is His or deleted;  
 $A_{33}$  is Thr or deleted;  
 $A_{34}$  is Ala or deleted;  
each of  $R_1$  and  $R_2$  is, independently, H,  $C_{1-12}$   
30 alkanyl,  $C_{7-20}$  phenylalkyl,  $C_{11-20}$  naphthylalkyl,  $C_{1-12}$ ,  
hydroxyalkyl,  $C_{2-12}$  hydroxyalkenyl,  $C_{7-20}$   
hydroxyphenylalkyl, or  $C_{11-20}$  hydroxynaphthylalkyl; or one  
and only one of  $R_1$  and  $R_2$  is  $COE_1$  in which  $E_1$  is  $C_{1-12}$   
alkyl,  $C_{2-12}$  alkyl,  $C_{2-12}$  alkenyl,  $C_{7-20}$  phenylalkyl,  $C_{11-20}$



- 31 -

napthylalkyl, C<sub>1-12</sub> hydroxyalkyl, C<sub>2-12</sub> hydroxyalkenyl, C<sub>7-20</sub> hydroxyphenylalkyl, or C<sub>11-20</sub> hydroxynapthylalkyl; and

R<sub>3</sub> is OH, NH<sub>2</sub>, C<sub>1-12</sub> alkoxy, or NH-Y-CH<sub>2</sub>-Z in which Y is a C<sub>1-12</sub> hydrocarbon moiety and Z is H, OH, CO<sub>2</sub>H or  
5 CONH<sub>2</sub>;

provided that at least one of A<sub>23</sub>, A<sub>24</sub>, A<sub>27</sub>, A<sub>28</sub>, or A<sub>31</sub> is Lys; or a pharmaceutically acceptable salt thereof.

29. A peptide of claim 28, wherein A<sub>22</sub> is Glu,  
10 Aib, or Cha; A<sub>23</sub> is Leu, Lys, or Cha; A<sub>25</sub> is Aib or Glu; A<sub>26</sub> is Aib or Lys; A<sub>28</sub> is Leu, Lys, or Cha; A<sub>29</sub> is Glu or Aib; A<sub>30</sub> is Cha, Aib, or Lys; A<sub>31</sub> is Leu, Cha, or Lys; A<sub>32</sub> is His; A<sub>33</sub> is Thr; and A<sub>34</sub> is Ala; or a pharmaceutically acceptable salt thereof.

15 30. A peptide of claim 29, wherein at least one of A<sub>7</sub> and A<sub>11</sub> is Cha; or a pharmaceutically acceptable salt thereof.

31. A peptide of claim 29, wherein at least one of A<sub>16</sub> or A<sub>19</sub> is Aib; or a pharmaceutically acceptable salt  
20 thereof.

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US96/11292

**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(6) : A61K 38/29; C07K 1/06, 14/635

US CL : 436/86; 530/324, 399; 514/12

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 436/86; 530/324, 399; 514/12

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, DIALOG

search terms: parathyroid hormone, parathyroid hormone related protein, PTH, PTHrP, derivatives, analogs, analogues

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHOREV et al. Modifications of Position 12 in Parathyroid Hormone and Parathyroid Hormone Related Protein: Toward the Design of Highly Potent Antagonists. Biochemistry. 1990, Vol. 29, No. 6, pages 1580-1586.	1, 8-11, 21, 28
Y	WO 94/02510 A2 (SANDOZ-ERFINDUNGEN VERWALTUNGSGELLSCHAFT) 03 February 1994, page 1, Abstract, page 74, lines 22 - 27.	1 and 8

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A* document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*E* earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
*O* document referring to an oral disclosure, use, exhibition or other means	
*P* document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

11 SEPTEMBER 1996

Date of mailing of the international search report

16 OCT 1996

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Authorized officer

KATHLEEN CARROLL

Facsimile No. (703) 305-3230

Telephone No. (703) 308-0196

Form PCT/ISA/210 (second sheet)(July 1992)\*